

AUTHOR INDEX

Bold type indicates authors of commentaries and editorials; (No.) indicates issue number.

- Agarwal JB, 49**
 Agnelli G, 9
 Albumin Reviewers, 6
Alcorn JM, 51
 Alderson P, 6, 66
Almbrand B, 5
 Anawalt B, 50
 Antonelli M, 10
 Avezum A, 7
- Badgett RG, 12, 74**
 Badrinath S, 50
Ball C, A-15-16(No.1)
Ballew KA, 44
 Barbone F, 48
Barer D, 59
 Barlow W, 42
 Barnett HJ, 33
Barry MJ, 61
Bates ER, 30
Bath P, 34
 Battié M, 42
Belknap SM, 39
 Benavente O, 59
 Bensoussan A, 74
Bergmann JF, 53
 Beyth RJ, 25
 Blum AL, 64
Boissel JP, 1
 Boshuizen HC, 20
 Boyd-Wickizer J, 17, 50
 Boyko EJ, 18
 Bradley KA, 17, 50
 Bredeson C, 53
 Broadhurst RJ, 77
 Broms K, 81
 Brooks MM, 49
 Browner WS, 78
 Bunn F, 6
 Buoncristiani P, 9
 Buring JE, 22
 Burman ML, 17
Burrows R, 47
 Bush K, 50
 Bush T, 8
- Calam J, 76**
 Campbell NC, 80
 Cantor SB, 43
Carey TS, 42
 Chalon S, 29
 Cherkin DC, 42
 Chevret S, 37
Choi P, 6
 Choudhri SH, 76
Clarfield AM, 20, 57
 Cline CM, 81
- Cochrane Injuries Group Albumin Reviewers, 6
 Colditz GA, 51
 Conti G, 10
Cook DJ, 6, 58
 Cortot A, 35
 Couper J, 40
 Curley C, 52
- Daurat V, 37
 Davey PG, 48
Davis MB, 20
 Degos F, 37
Delaney BC, 19
 Deyo RA, 42
 Diamond S, 12
Diemer M, 22
 Dodick DW, 11
 Douglas PS, 45
- Eckman MH, 24
 Edmeads JG, 11
 Eliasziw M, 33
 Elkind A, 12
Elliott CG, 75
 El-Omar E, 65
 Embil JM, 76
Engel CC Jr, 43
 Erhardt LR, 81
 European Study Group on Interferon β -1b in Secondary Progressive MS, 69
 Ewe K, 36
- Falk RH, 24
 Färkkilä MA, 71
Feagan BG, 36
Feinberg AW, 18
Fekete T, 15
Ferguson R, 43
 Fieschi C, 34
 Fleischmann KE, 45
Fletcher RH, 13
 Forette F, 57
- Garg PP, 23
 Garner P, 15
Geerts WH, 9
Gerstein HC, 2-3
 Ginsberg JS, 75
 Giovannucci E, 51
 Girardi LS, 63
 Glasziou P, 13
Goodman S, 46
 Grady D, 8, 23
Green MS, 24
 Griffiths AM, 36
Gurwitz JH, 48
 Guyatt GH, 44
- Hacke W, 34
 Hakala K, 71
 Hankey GJ, 77
 Hanna SE, 79
 Hauth J, 47
Hawkey CJ, 38, 64-65
 Haynes RB, 79
 Hemmelgarn BR, 44
 Henderson RA, 30
 Hennekens CH, 22
Herr DL, 66
Hinchey JA, 62
 Hing M, 74
Hirsch CH, 67
 Holloway RG, 62
 Holzer CE III, 43
 Hopkins RJ, 63
Hotopf M, 60
 House JS, 21
Hughes CM, 58
 Hulley S, 8
Humphrey LL, 21
 Hunink MG, 45
 Hunt DL, 79
- International Hepatitis Interventional Therapy (IHIT) Group, 70
Irvine EJ, 35
 Irwig L, 13
 Ishani A, 61
 Israelsson BY, 81
 Izaks GJ, 20
- Jackson RT, 12
 Jacobs AK, 49
Jacobs M, 45
 Jamrozik K, 77
 Jewell D, 35
Jones JIW, 64-65
 Jurriaans S, 14
- Kaste M, 34
Kauhanen J, 50
 Kearon C, 75
Keenan SP, 10, 58
 Kelsey SF, 49
 Kerlikowske K, 23
 Kewenter J, 13
King PD, 70
Korula J, 37
 Kuntz KM, 45
- Lader E, 7**
 Laine L, 63
 Lancaster T, 41
 Landefeld S, 25
Langer RD, 8

- Lantz PM, 21
Laupacis A, 72-73
 Lechat P, 29
 Lee SS, 70
 Lefebvre C, 6
 Lepkowski JM, 21
 Lewis SJ, 32
 Ligthart GJ, 20
 Linde K, 60
 Lindheimer M, 47
 Liu M, 62
Longstreth W, 33
 The Long-Term Intervention with
 Pravastatin in Ischaemic Disease
 (LIPID) Study Group, 31
Luxenberg JS, 78
- Mackey S, 16
Malmberg K, 5
 Marcellin P, 70
 Marrugat J, 46
 Masia R, 46
 Matias M, 22
McAlister FA, A-13(No. 3)
 McColl K, 65
McDonald JWD, 71
 McEachern JE, 52
 McGee SR, 18
 McMahon AD, 48
Menken M, 11
 Meyers PD, 39
Mills JK, 80
 Modigliani R, 36
 Moher D, 59
 Moye LA, 32
 Mulrow CD, 60
 Multicentre GER-CYT-04 Group, 37
 Murray L, 65
- Nandurkar S, 19
 National Institute of Child Health and
 Human Development Network of
 Maternal-Fetal Medicine Units, 47
 Nichol G, 24
 North American Symptomatic Carotid
 Endarterectomy Trial Collaborators, 33
- O'Connor P, 68-69
 O'Morain C, 64
 Omeprazole plus Clarithromycin and
 Amoxicillin Effect One Year after
 Treatment (OCAY) Study Group, 64
 Osteoporotic Fractures Research Study
 Group, 78
- Packer M, 29
 Palmer M, 67
- Paltiel O, 57
 Pang D, 58
 Panju AA, 44
 Pauker SG, 24
Peterson WL, 25, 63
 Pham B, 59
 Piovela F, 9
 Pocock SJ, 30
 Pogue J, 7
 Powell SH, 17
 Poynard T, 70
 Price JR, 40
 PRISMS (Prevention of Relapses
 and Disability by Interferon β -1a
 Subcutaneously in Multiple Sclerosis)
 Study Group, 69
 Pryse-Phillips WE, 11
Pulicino P, 77
 The PURSUIT Trial Investigators, 1
- Quinn LM, 25
- Randomised Intervention Treatment of
 Angina (RITA-1) trial participants, 30
 Reijers MH, 14
 Ridker PM, 22
 Ritchie LD, 80
 Roberts I, 66
 Rocco M, 10
 Rodger M, 53
 Roe B, 67
Rubin HR, 52
Rydén L, 5
- Sacks FM, 32
Saitz R, 17
 Sala J, 46
 Schierhout G, 66
Schmedtje JF Jr, 29
 Second European-Australasian Acute
 Stroke Study Investigators, 34
Selby R, 9
 Seux ML, 57
 Sharp SJ, 30
 Sheehan M, 16
Sherman SE, 16
 Shih J, 22
 Sibai BM, 47
 Sibbald WJ, 58
 Simel DL, 44
 Sirinavin S, 15
 Smart G, 76
 Smith B, 79
Smith DE, 14
 Smith K, 79
 Speroff T, 52
 Staessen JA, 57
- Stampfer MJ, 51
 Stark G, 61
 Stead LF, 41
 Stefanick ML, 16
 Steinbauer JR, 43
 Steinhart AH, 36
Strauss SE, A-13(No. 3)
 Street J, 42
 Study of Osteoporotic Fractures
 Research Group, 78
 Subak L, 23
 Syst-Eur Investigators, 57
 Szczepanski L, 38
- Taha AS, 38**
 Talley NJ, 19, 64, 74
 Taylor DW, 33
Taylor WC, 31
Thacker HL, 23
 Thain J, 80
 Thomsen OO, 35, 36
 Towler BP, 13
 Treiman DM, 39
 Tsuyuki RT, 7
 Tulassay Z, 38
 Turunen UM, 71
- UK Prospective Diabetes
 Study Group, 2-3, 4-5
 UK Small Aneurysm Trial
 Participants, 72-73
- van Buuren S, 20
 Veterans Affairs Status Epilepticus
 Cooperative Study Group, 39
 Vickers A, A-13-14(No. 2)
 Volk RJ, 43
- Walton NY, 39
 Wardlaw J, 62
Weekley J, 79
 Weller D, 13
 Wells PS, 53, 75
Wessely S, 40
Westman EC, 41
 Weverling GJ, 14
 Whooley MA, 78
 Willenheimer RB, 81
 Williams K, 67
 Wilt TJ, 61
Woods KL, 32
- Xia H, 19
- Yusuf S, 7

SUBJECT INDEX

A

- Abruptio placentae
 - Preeclampsia and baseline proteinuria in women with chronic hypertension were associated with adverse neonatal outcomes, 47
- Adrenal cortex hormones
 - Review: Insufficient evidence exists for the effectiveness of 5 commonly used interventions for severe head injury, 66
- Adrenergic β -antagonists
 - Review: β -blockers reduce mortality and morbidity in congestive heart failure, 7
 - Review: β -blockers benefit patients with chronic heart failure, 29
- Aged
 - Advanced age predicted risk for recurrent stroke after first-ever stroke, 77
- Antihypertensive treatment may have reduced the rate of dementia in older patients with isolated systolic hypertension, 57
- Depressive symptoms were associated with increased mortality in older women, 78
- Low blood pressure was not associated with increased mortality in very elderly persons after adjustment for poor health status, 20
- Pravastatin reduced cardiovascular events in older patients with myocardial infarction and average cholesterol levels, 32
- Albumin
 - Review: Albumin does not reduce death in critically ill patients with hypovolemia, burns, or hypoalbuminemia, 6
- Alcohol
 - The Alcohol Use Disorders Identification Test was not affected by ethnic and sex bias, 43
 - Low income was an independent risk factor for premature death after controlling for health behaviors, 21
 - Review: Some alcohol screening tests have acceptable test properties for use in general clinical populations of U.S. women, 17
 - Review: Some health risks associated with drinking alcohol are greater for women than for men, 50
- Alzheimer disease
 - Antihypertensive treatment may have reduced the rate of dementia in older patients with isolated systolic hypertension, 57
- Amoxicillin
 - Eradication of *Helicobacter pylori* did not relieve symptoms of nonulcer dyspepsia but increased healing of gastritis, 64
 - Eradication of *Helicobacter pylori* with omeprazole and antibiotics improved symptoms in nonulcer dyspepsia, 65
 - Review: Successful *Helicobacter pylori* eradication prevents ulcer recurrence, 63
- Angina pectoris
 - Eptifibatide reduced deaths and nonfatal myocardial infarction in acute coronary syndromes, 1
 - PTCA and CABG had similar long-term mortality and myocardial infarction rates, 30
- Angioplasty, transluminal, percutaneous coronary
 - Women had a lower risk than men for mortality after coronary revascularization, 49
- Antiarrhythmia agents
 - Cardioversion followed by aspirin alone or with amiodarone was cost-effective for nonvalvular atrial fibrillation, 24
- Antibiotics
 - Eradication of *Helicobacter pylori* did not relieve symptoms of nonulcer dyspepsia but increased healing of gastritis, 64
 - Eradication of *Helicobacter pylori* with omeprazole and antibiotics improved symptoms in nonulcer dyspepsia, 65
 - Review: Antibiotics have no effect on diarrhea and increase bacteriologic relapse in salmonella intestinal infection, 15
- Anticholesteremic agents
 - Pravastatin reduced cardiovascular events in older patients with myocardial infarction and average cholesterol levels, 32
- Anticoagulants
 - A 5-point scoring system predicted bleeding in patients treated with warfarin, 25
- Anticonvulsants
 - 4 anticonvulsant drug regimens had similar efficacies for generalized convulsive status epilepticus, 39
- Antidepressants
 - Review: St. John's wort is more effective than placebo for treating depressive disorders, 60
- Anti-HIV agents
 - Dual maintenance therapy did not sustain suppression of viral replication in HIV-1 infection compared with prolonged reduction therapy, 14
- Antihypertensive agents
 - Antihypertensive treatment may have reduced the rate of dementia in older patients with isolated systolic hypertension, 57
 - Tight blood pressure control reduced diabetes mellitus-related deaths and complications and was cost-effective in type 2 diabetes, 4-5
- Anti-inflammatory agents
 - Budesonide increased remission rates more than mesalamine in active Crohn disease, 35
 - Helicobacter pylori* eradication did not change ulcer recurrence or dyspepsia rates in patients who used NSAIDs, 38
- Aortic aneurysm, abdominal
 - Early surgery was more expensive than surveillance for small abdominal aortic aneurysms but slightly improved health perception and bodily pain, 72
 - Early surgery did not reduce 6-year mortality in patients with small abdominal aortic aneurysms, 73
- Arterial occlusive diseases
 - Review: Physical examination predicts the presence and distribution of peripheral vascular disease, 18
- Aspirin
 - Cardioversion followed by aspirin alone or with amiodarone was cost-effective for nonvalvular atrial fibrillation, 24
 - Helicobacter pylori* infection was not associated with nonulcer dyspepsia, 19
- Atrial fibrillation
 - Cardioversion followed by aspirin alone or with amiodarone was cost-effective for nonvalvular atrial fibrillation, 24

B

- Behavior therapy
 - Review: Inconclusive evidence suggests that bladder training improves urge incontinence, 67
- Benzodiazepine
 - Anxiolytic benzodiazepines and zopiclone increased road traffic crashes, 48
- Blood glucose
 - Intensive blood glucose control reduced type 2 diabetes mellitus-related end points, 2
- Blood pressure
 - Low blood pressure was not associated with increased mortality in very elderly persons after adjustment for poor health status, 20
 - Tight blood pressure control reduced diabetes mellitus-related deaths and complications and was cost-effective in type 2 diabetes, 4-5
- Blood proteins
 - Review: Albumin does not reduce death in critically ill patients with hypovolemia, burns, or hypoalbuminemia, 6
- Body weight
 - Low income was an independent risk factor for premature death after controlling for health behaviors, 21
- Brain injuries
 - Review: Insufficient evidence exists for the effectiveness of 5 commonly used interventions for severe head injury, 66
- Breast neoplasms
 - Review: Some health risks associated with drinking alcohol are greater for women than for men, 50

C

- Cardiovascular disease
 - C-reactive protein levels were associated with cardiovascular events in apparently healthy women, 22
 - Pravastatin reduced all-cause mortality, CHD mortality, and cardiovascular disease mortality in patients with existing CHD, 31
 - Pravastatin reduced cardiovascular events in older patients with myocardial infarction and average cholesterol levels, 32
 - Review: Some health risks associated with drinking alcohol are greater for women than for men, 50
- Carotid stenosis
 - Carotid endarterectomy had a modest benefit for 50% to 69% stenosis after TIA or nondisabling stroke and no benefit for < 50% stenosis, 33
 - Review: Carotid endarterectomy modestly reduces ipsilateral stroke in asymptomatic stenosis, 59
- Cerebral hemorrhage
 - Advanced age predicted risk for recurrent stroke after first-ever stroke, 77
- Cerebral ischemia
 - Alteplase given within 6 hours of stroke did not improve clinical outcomes at 90 days, 34
 - Carotid endarterectomy had a modest benefit for 50% to 69% stenosis after TIA or nondisabling stroke and no benefit for < 50% stenosis, 33

- Cerebrovascular disorders
 - Advanced age predicted risk for recurrent stroke after first-ever stroke, 77
 - Carotid endarterectomy had a modest benefit for 50% to 69% stenosis after TIA or nondisabling stroke and no benefit for < 50% stenosis, 33
 - Review: Carotid endarterectomy modestly reduces ipsilateral stroke in asymptomatic stenosis, 59
 - Review: Data are insufficient to evaluate the effectiveness of doses and administration routes of thrombolytic agents for acute ischemic stroke, 62
- Chiropractic
 - Chiropractic manipulation and McKenzie physiotherapy were not effective for low back pain, 42
- Cholesterol
 - Pravastatin reduced cardiovascular events in older patients with myocardial infarction and average cholesterol levels, 32
- Ciprofloxacin
 - Ciprofloxacin reduced treatment failure in ulcerative colitis at 6 months but not at 12 months, 71
- Clarithromycin
 - Eradication of *Helicobacter pylori* did not relieve symptoms of nonulcer dyspepsia but increased healing of gastritis, 64
 - Review: Successful *Helicobacter pylori* eradication prevents ulcer recurrence, 63
- Cognitive therapy
 - Review: Cognitive behavior therapy is effective for the chronic fatigue syndrome, 40
- Colitis, ulcerative
 - Ciprofloxacin reduced treatment failure in ulcerative colitis at 6 months but not at 12 months, 71
- Colloids
 - Review: Albumin does not reduce death in critically ill patients with hypovolemia, burns, or hypoalbuminemia, 6
- Colonic neoplasms
 - Long-term dietary folate was associated with a decreased risk for colon cancer in women, 51
- Colorectal neoplasms
 - Review: Fecal occult blood test screening reduces colorectal cancer mortality, 13
- Congestive heart failure
 - Nurse-run clinics in primary care increased secondary prevention in coronary artery disease, 80
 - A nurse-run program for congestive heart failure reduced time to hospital readmission, 81
 - Review: β -blockers benefit patients with chronic heart failure, 29
 - Review: β -blockers reduce mortality and morbidity in congestive heart failure, 7
- Coronary artery bypass graft
 - PTCA and CABG had similar long-term mortality and myocardial infarction rates, 30
 - Women had a lower risk than men for mortality after coronary revascularization, 49
- Coronary disease
 - Estrogen plus progestin was not effective for secondary prevention of coronary heart disease in postmenopausal women, 8

- Nurse-run clinics in primary care increased secondary prevention in coronary heart disease, 80
- A nurse-run program for congestive heart failure reduced time to hospital readmission, 81
- PTCA and CABG had similar long-term mortality and myocardial infarction rates, 30
- Pravastatin reduced all-cause mortality, CHD mortality, and cardiovascular disease mortality in patients with existing CHD, 31
- Review: Exercise ECHO is a more specific and discriminatory test than exercise SPECT for coronary artery disease, 45
- Women had a lower risk than men for mortality after coronary revascularization, 49
- Costs and cost analysis
 - Nurse-run clinics in primary care increased secondary prevention in coronary artery disease, 80
 - A nurse-run program for congestive heart failure reduced time to hospital readmission, 81
 - Tight blood pressure control reduced diabetes mellitus-related deaths and complications and was cost-effective in type 2 diabetes, 4-5
- C-reactive protein
 - C-reactive protein levels were associated with cardiovascular events in apparently healthy women, 22
- Critical illness
 - Review: Albumin does not reduce death in critically ill patients with hypovolemia, burns, or hypoalbuminemia, 6
- Crohn disease
 - Budesonide increased remission rates more than mesalamine in active Crohn disease, 35
 - Review: Corticosteroid therapy does not reduce the rate of relapse in Crohn disease, 36

D

- D-dimer assay
 - D-dimer assay was useful for ruling out suspected pulmonary embolism, 75
- Decision support systems, clinical
 - Review: Computer-based clinical decision support systems can improve physician performance in some areas, 79
- Dementia
 - Antihypertensive treatment may have reduced the rate of dementia in older patients with isolated systolic hypertension, 57
- Depression
 - Depressive symptoms were associated with increased mortality in older women, 78
 - Review: St. John's wort is more effective than placebo for treating depressive disorders, 60
- Diabetes mellitus
 - Intensive blood glucose control reduced type 2 diabetes mellitus-related end points, 2
 - Metformin reduced diabetes-related end points and all-cause mortality in overweight patients with type 2 diabetes, 3

- Tight blood pressure control reduced diabetes mellitus-related deaths and complications and was cost-effective in type 2 diabetes, 4-5
- Diarrhea
 - Review: Antibiotics have no effect on diarrhea and increase bacteriologic relapse in salmonella intestinal infection, 15
- Diet
 - Exercise and diet improved cholesterol levels in high-risk adults, 16
 - Long-term dietary folate was associated with a decreased risk for colon cancer in women, 51
- Direct service costs
 - Early surgery was more expensive than surveillance for small abdominal aortic aneurysms but slightly improved health perception and bodily pain, 72
- Diuretics, osmotic
 - Review: Insufficient evidence exists for the effectiveness of 5 commonly used interventions for severe head injury, 66
- Drug therapy
 - Dual maintenance therapy did not sustain suppression of viral replication in HIV-1 infection compared with prolonged reduction therapy, 14
 - Intensive blood glucose control reduced type 2 diabetes mellitus-related end points, 2
 - Metformin reduced diabetes mellitus-related end points and all-cause mortality in overweight patients with type 2 diabetes, 3
 - Review: β -blockers reduce mortality and morbidity in congestive heart failure, 7
- Duodenal ulcer
 - Review: Successful *Helicobacter pylori* eradication prevents ulcer recurrence, 63
- Dyspepsia
 - Eradication of *Helicobacter pylori* did not relieve symptoms of nonulcer dyspepsia but increased healing of gastritis, 64
 - Eradication of *Helicobacter pylori* with omeprazole and antibiotics improved symptoms in nonulcer dyspepsia, 65
 - Helicobacter pylori* eradication did not change ulcer recurrence or dyspepsia rates in patients who used NSAIDs, 38
 - Helicobacter pylori* infection was not associated with nonulcer dyspepsia, 19

E

- Echocardiography
 - Review: Exercise ECHO is a more specific and discriminatory test than exercise SPECT for coronary artery disease, 45
- Editorial
 - A clinician's guide to journal articles about prognosis, A-13(No. 3)
 - Evidence-based medicine and complementary medicine, A-13(No. 2)
 - Evidence-based medicine on the wards: report from an evidence-based minion, A-15(No.1)

Electrocardiography

Review: Clinical features and electrocardiographic changes predict myocardial infarction in adults with chest pain, 44

Endarterectomy, carotid

Carotid endarterectomy had a modest benefit for 50% to 69% stenosis after TIA or nondisabling stroke and no benefit for < 50% stenosis, 33

Review: Carotid endarterectomy modestly reduces ipsilateral stroke in asymptomatic stenosis, 59

Endoscopy

A negative serum enzyme immunoassay test result ruled out *Helicobacter pylori* infection, 76

Enoxaparin

Enoxaparin reduced venous thromboembolism after elective neurosurgery, 9

Eptifibatide

Eptifibatide reduced deaths and nonfatal myocardial infarction in acute coronary syndrome, 1

Estrogen replacement therapy

Estrogen plus progestin was not effective for secondary prevention of coronary heart disease in postmenopausal women, 8

Review: Postmenopausal hormone replacement therapy is associated with increased invasive ovarian cancer, especially after long-term use, 23

Exercise

Exercise and diet improved cholesterol levels in high-risk adults, 16

Low income was an independent risk factor for premature death after controlling for health behaviors, 21

Review: Exercise ECHO is a more specific and discriminatory test than exercise SPECT for coronary artery disease, 45

F

Fatigue syndrome, chronic

Review: Cognitive behavior therapy is effective for the chronic fatigue syndrome, 40

Fibrin fibrinogen degradation products

D-dimer assay was useful for ruling out suspected pulmonary embolism, 75

Fibrinolytic agents

Alteplase given within 6 hours of stroke did not improve clinical outcomes at 90 days, 34

Cardioversion followed by aspirin alone or with amiodarone was cost-effective for nonvalvular atrial fibrillation, 24

Folic acid

Long-term dietary folate was associated with a decreased risk for colon cancer in women, 51

Function

Review: Cognitive behavior therapy is effective for the chronic fatigue syndrome, 40

G

Gastroenteritis

Review: Antibiotics have no effect on diarrhea and increase bacteriologic relapse in salmonella intestinal infection, 15

Glucocorticoids

Budesonide increased remission rates more than mesalamine in active Crohn disease, 35

Review: Corticosteroid therapy does not reduce the rate of relapse in Crohn disease, 36

H

Health behavior

Low income was an independent risk factor for premature death after controlling for health behaviors, 21

Health status

Low blood pressure was not associated with increased mortality in very elderly persons after adjustment for poor health status, 20

Helicobacter pylori

Eradication of *Helicobacter pylori* did not relieve symptoms of nonulcer dyspepsia but increased healing of gastritis, 64

Eradication of *Helicobacter pylori* with omeprazole and antibiotics improved symptoms in nonulcer dyspepsia, 65

Helicobacter pylori eradication did not change ulcer recurrence or dyspepsia rates in patients who used NSAIDs, 38

Helicobacter pylori infection was not associated with nonulcer dyspepsia, 19

A negative serum enzyme immunoassay test result ruled out *Helicobacter pylori* infection, 76

Review: Successful *Helicobacter pylori* eradication prevents ulcer recurrence, 63

Hemorrhage

A 5-point scoring system predicted bleeding in patients treated with warfarin, 25

Heparin

Low-molecular-weight heparin was more cost-effective than unfractionated heparin for treating deep venous thrombosis in several clinical settings, 53

Heparin, low-molecular-weight

Low-molecular-weight heparin was more cost-effective than unfractionated heparin for treating deep venous thrombosis in several clinical settings, 53

Hepatitis C

A higher dose and longer duration of interferon α -2a reduced the incidence of cirrhosis in chronic hepatitis C, 37

Interferon- α 2b and ribavirin increased the loss of HCV RNA in chronic hepatitis C infection, 70

HIV infections

Dual maintenance therapy did not sustain suppression of viral replication in HIV-1 infection compared with prolonged reduction therapy, 14

- Hospital charges
Interdisciplinary medical rounds improved patient care efficiency and the care process and decreased costs, 52
- Hospitalization
Review: β -blockers benefit patients with chronic heart failure, 29
- Hypertension
Antihypertensive treatment may have reduced the rate of dementia in older patients with isolated systolic hypertension, 57
Preeclampsia and baseline proteinuria in women with chronic hypertension were associated with adverse neonatal outcomes, 47
Tight blood pressure control reduced diabetes mellitus-related deaths and complications and was cost-effective in type 2 diabetes, 4-5
- Hypoglycemic agents
Intensive blood glucose control reduced type 2 diabetes mellitus-related end points, 2
Metformin reduced diabetes mellitus-related end points and all-cause mortality in overweight patients with type 2 diabetes, 3

I

- Immunoassay
A negative serum enzyme immunoassay test result ruled out *Helicobacter pylori* infection, 76
- Incontinence
Review: Inconclusive evidence suggests that bladder training improves urge incontinence, 67
- Inflammatory bowel disease
Review: Corticosteroid therapy does not reduce the rate of relapse in Crohn disease, 36
- Insulin
Intensive blood glucose control reduced type 2 diabetes mellitus-related end points, 2
- Intensive care units
Noninvasive positive-pressure ventilation was as effective as conventional mechanical ventilation and had fewer complications, 10
- Interferons
A higher dose and longer duration of interferon α -2a reduced the incidence of cirrhosis in chronic hepatitis C, 37
Interferon- α 2b and ribavirin increased the loss of HCV RNA in chronic hepatitis C infection, 70
Interferon- β 1a reduced relapses at 2 years in relapsing-remitting multiple sclerosis, 68
Interferon- β 1b reduced the progression of secondary progressive multiple sclerosis, 69
- Interprofessional relations
Interdisciplinary medical rounds improved patient care efficiency and the care process and decreased costs, 52
- Intranasal administration
Sumatriptan nasal spray was effective and well tolerated for multiple migraine attacks, 12
- Intubation, intratracheal
Noninvasive positive-pressure ventilation was as effective as conventional mechanical ventilation and had fewer complications, 10

- Review: Continuous positive airway pressure support reduces the need for intubation in patients with cardiogenic pulmonary edema, 58
- Irritable bowel syndrome
Chinese herbal medicine was effective in the irritable bowel syndrome, 74
- Ischemia
Review: Physical examination predicts the presence and distribution of peripheral vascular disease, 18

L

- Lipids
Exercise and diet improved cholesterol levels in high-risk adults, 16
Pravastatin reduced all-cause mortality, CHD mortality, and cardiovascular disease mortality in patients with existing CHD, 31
- Liver cirrhosis
A higher dose and longer duration of interferon α -2a reduced the incidence of cirrhosis in chronic hepatitis C, 37
- Low back pain
Chiropractic manipulation and McKenzie physiotherapy were not effective for low back pain, 42

M

- Mannitol
Review: Insufficient evidence exists for the effectiveness of 5 commonly used interventions for severe head injury, 66
- Medical history taking
Review: Clinical features and electrocardiographic changes predict myocardial infarction in adults with chest pain, 44
- Medicine, herbal
Chinese herbal medicine was effective in the irritable bowel syndrome, 74
Review: St. John's wort is more effective than placebo for treating depressive disorders, 60
- Menopause
Review: Postmenopausal hormone replacement therapy is associated with increased invasive ovarian cancer, especially after long-term use, 23
- Mesalamine
Budesonide increased remission rates more than mesalamine in active Crohn disease, 35
- Metformin
Metformin reduced diabetes mellitus-related end points and all-cause mortality in overweight patients with type 2 diabetes, 3
- Metronidazole
Eradication of *Helicobacter pylori* with omeprazole and antibiotics improved symptoms in nonulcer dyspepsia, 65
- Migraine
Review: Several nonpharmacologic treatments are effective for migraine, especially when used with other agents or for prophylaxis, 11
Sumatriptan nasal spray was effective and well tolerated for multiple migraine attacks, 12

Mortality

- Depressive symptoms were associated with increased mortality in older women, 78
- Eptifibatide reduced deaths and nonfatal myocardial infarction in acute coronary syndromes, 1
- Low blood pressure was not associated with increased mortality in very elderly persons after adjustment for poor health status, 20
- Low income was an independent risk factor for premature death after controlling for health behaviors, 21
- Review: β -blockers benefit patients with chronic heart failure, 29
- Review: Some health risks associated with drinking alcohol are greater for women than for men, 50
- Women had a worse prognosis than men after myocardial infarction, 46

Multiple sclerosis

- Interferon- β 1a reduced relapses at 2 years in relapsing-remitting multiple sclerosis, 68
- Interferon- β 1b reduced the progression of secondary progressive multiple sclerosis, 69

Myocardial infarction

- Eptifibatide reduced deaths and nonfatal myocardial infarction in acute coronary syndromes, 1
- Pravastatin reduced all-cause mortality, CHD mortality, and cardiovascular disease mortality in patients with existing CHD, 31
- PTCA and CABG had similar long-term mortality and myocardial infarction rates, 30
- Review: Clinical features and electrocardiographic changes predict myocardial infarction in adults with chest pain, 44
- Women had a lower risk than men for mortality after coronary revascularization, 49
- Women had a worse prognosis than men after myocardial infarction, 46

N

Neonatal outcomes

- Preeclampsia and baseline proteinuria in women with chronic hypertension were associated with adverse neonatal outcomes, 47

Neurosurgery

- Enoxaparin reduced venous thromboembolism after elective neurosurgery, 9

Nurse clinicians

- Nurse-run clinics in primary care increased secondary prevention in coronary heart disease, 80
- A nurse-run program for congestive heart failure reduced time to hospital readmission, 81

O

Obesity, in diabetes

- Metformin reduced diabetes mellitus-related end points and all-cause mortality in overweight patients with type 2 diabetes, 3

Occult blood

- Review: Fecal occult blood test screening reduces colorectal cancer mortality, 13

Omeprazole

- Eradication of *Helicobacter pylori* did not relieve symptoms of nonulcer dyspepsia but increased healing of gastritis, 64
- Eradication of *Helicobacter pylori* with omeprazole and antibiotics improved symptoms in nonulcer dyspepsia, 65
- Review: Successful *Helicobacter pylori* eradication prevents ulcer recurrence, 63

Organization and administration

- Interdisciplinary medical rounds improved patient care efficiency and the care process and decreased costs, 52

Outcome and process assessment

- Review: Computer-based clinical decision support systems can improve physician performance in some areas, 79

Ovarian neoplasms

- Review: Postmenopausal hormone replacement therapy is associated with increased invasive ovarian cancer, especially after long-term use, 23

P

Pain

- Review: Clinical features and electrocardiographic changes predict myocardial infarction in adults with chest pain, 44

Patient care planning

- Nurse-run clinics in primary care increased secondary prevention in coronary artery disease, 80
- A nurse-run program for congestive heart failure reduced time to hospital readmission, 81

Patient education

- Chiropractic manipulation and McKenzie physiotherapy were not effective for low back pain, 42
- Nurse-run clinics in primary care increased secondary prevention in coronary artery disease, 80
- A nurse-run program for congestive heart failure reduced time to hospital readmission, 81

Patient satisfaction

- Interdisciplinary medical rounds improved patient care efficiency and staff satisfaction and decreased costs, 52

Peptic ulcer

- Helicobacter pylori* eradication did not change ulcer recurrence or dyspepsia rates in patients who used NSAIDs, 38

Peptides

- Eptifibatide reduced deaths and nonfatal myocardial infarction in acute coronary syndromes, 1

Percutaneous transluminal coronary angioplasty

- PTCA and CABG had similar long-term mortality and myocardial infarction rates, 30

Physical examination

- Review: Clinical features and electrocardiographic changes predict myocardial infarction in adults with chest pain, 44
- Review: Physical examination predicts the presence and distribution of peripheral vascular disease, 18

Physical therapy

Chiropractic manipulation and McKenzie physiotherapy were not effective for low back pain, 42

Plants, medicinal

Chinese herbal medicine was effective in the irritable bowel syndrome, 74

Review: Saw palmetto plant extract (*Serenoa repens*) is effective and safe in benign prostatic hyperplasia, 61

Review: St. John's wort is more effective than placebo for treating depressive disorders, 60

Platelet aggregation inhibitors

Eptifibatide reduced deaths and nonfatal myocardial infarction in acute coronary syndromes, 1

Positive-pressure ventilation

Noninvasive positive-pressure ventilation was as effective as conventional mechanical ventilation and had fewer complications, 10

Review: Continuous positive airway pressure support reduces the need for intubation in patients with cardiogenic pulmonary edema, 58

Postmenopause

Estrogen plus progestin was not effective for secondary prevention of coronary heart disease in postmenopausal women, 8

Exercise and diet improved cholesterol levels in high-risk adults, 16

Review: Postmenopausal hormone replacement therapy is associated with increased ovarian cancer, especially after long-term use, 23

Pravastatin

Pravastatin reduced all-cause mortality, CHD mortality, and cardiovascular disease mortality in patients with existing CHD, 31

Pravastatin reduced cardiovascular events in older patients with myocardial infarction and average cholesterol levels, 32

Pregnancy

Preeclampsia and baseline proteinuria in women with chronic hypertension were associated with adverse neonatal outcomes, 47

Primary health care

The Alcohol Use Disorders Identification Test was not affected by ethnic and sex bias, 43

Progestational hormones

Estrogen plus progestin was not effective for secondary prevention of coronary heart disease in postmenopausal women, 8

Prostatic hyperplasia

Review: Saw palmetto plant extract (*Serenoa repens*) is effective and safe in benign prostatic hyperplasia, 61

Psychotherapy

Review: Several nonpharmacologic treatments are effective for migraine, especially when used with other agents or for prophylaxis, 11

Psychotropic drugs

Anxiolytic benzodiazepines and zopiclone increased road traffic crashes, 48

Pulmonary edema

Review: Continuous positive airway pressure support reduces the need for intubation in patients with cardiogenic pulmonary edema, 58

Pulmonary embolism

D-dimer assay was useful for ruling out suspected pulmonary embolism, 75

Enoxaparin reduced venous thromboembolism after elective neurosurgery, 9

Q

Quality of life

Early surgery was more expensive than surveillance for small abdominal aortic aneurysms but slightly improved health perception and bodily pain, 72

Review: Cognitive behavior therapy is effective for the chronic fatigue syndrome, 40

Questionnaires

The Alcohol Use Disorders Identification Test was not affected by ethnic and sex bias, 43

Review: Some alcohol screening tests have acceptable test properties for use in general clinical populations of U.S. women, 17

R

Ranitidine

Review: Successful *Helicobacter pylori* eradication prevents ulcer recurrence, 63

Recurrence

Interferon- β 1a reduced relapses at 2 years in relapsing-remitting multiple sclerosis, 68

Interferon- β 1b reduced the progression of secondary progressive multiple sclerosis, 69

Review: Successful *Helicobacter pylori* eradication prevents ulcer recurrence, 63

Respiration, artificial

Noninvasive positive-pressure ventilation was as effective as conventional mechanical ventilation and had fewer complications, 10

Retreatment

Ciprofloxacin reduced treatment failure in ulcerative colitis at 6 months but not at 12 months, 71

Ribavirin

Interferon- α 2b and ribavirin increased the loss of HCV RNA in chronic hepatitis C infection, 70

S

Salmonella infections

Review: Antibiotics have no effect on diarrhea and increase bacteriologic relapse in salmonella intestinal infection, 15

Sedatives, barbiturate

Review: Insufficient evidence exists for the effectiveness of 5 commonly used interventions for severe head injury, 66

Sestamibi

Review: Exercise ECHO is a more specific and discriminatory test than exercise SPECT for coronary artery disease, 45

Sex factors

Review: Some health risks associated with drinking alcohol are greater for women than for men, 50

Women had a lower risk than men for mortality after coronary revascularization, 49

Women had a worse prognosis than men after myocardial infarction, 46

Shock, cardiogenic

Review: Continuous positive airway pressure support reduces the need for intubation in patients with cardiogenic pulmonary edema, 58

Smoking

Helicobacter pylori infection was not associated with nonulcer dyspepsia, 19

Low income was an independent risk factor for premature death after controlling for health behaviors, 21

Review: Self-help interventions alone minimally increase smoking cessation, 41

Socioeconomic factors

Low income was an independent risk factor for premature death after controlling for health behaviors, 21

Status epilepticus

4 anticonvulsant drug regimens had similar efficacies for generalized convulsive status epilepticus, 39

Sumatriptan

Sumatriptan nasal spray was effective and well tolerated for multiple migraine attacks, 12

Surgical procedures, elective

Early surgery was more expensive than surveillance for small abdominal aortic aneurysms but slightly improved health perception and bodily pain, 72

Early surgery did not reduce 6-year mortality in patients with small abdominal aortic aneurysms, 73

T

Tetracycline

Eradication of *Helicobacter pylori* with omeprazole and antibiotics improved symptoms in nonulcer dyspepsia, 65

Thallium

Review: Exercise ECHO is a more specific and discriminatory test than exercise SPECT for coronary artery disease, 45

Thromboembolism

Enoxaparin reduced venous thromboembolism after elective neurosurgery, 9

Thrombolytic therapy

Alteplase given within 6 hours of stroke did not improve clinical outcomes at 90 days, 34

Review: Data are insufficient to evaluate the effectiveness of doses and administration routes of thrombolytic agents for acute ischemic stroke, 62

Tissue plasminogen activator

Alteplase given within 6 hours of stroke did not improve clinical outcomes at 90 days, 34

Traffic accidents

Anxiolytic benzodiazepines and zopiclone increased road traffic crashes, 48

U

Urinary incontinence

Review: Inconclusive evidence suggests that bladder training improves urge incontinence, 67

V

Vascular disease

Review: Physical examination predicts the presence and distribution of peripheral vascular disease, 18

Vasoconstrictor agents

Sumatriptan nasal spray was effective and well tolerated for multiple migraine attacks, 12

Venous thrombosis

Low-molecular-weight heparin was more cost-effective than unfractionated heparin for treating deep venous thrombosis in several clinical settings, 53

Ventilation-perfusion ratio

D-dimer assay was useful for ruling out suspected pulmonary embolism, 75

Ventricular function, left

Review: β -blockers benefit patients with chronic heart failure, 29

Viral RNA

Dual maintenance therapy did not sustain suppression of viral replication in HIV-1 infection compared with prolonged reduction therapy, 14

Vitamins

Long-term dietary folate was associated with a decreased risk for colon cancer in women, 51

W

Women

C-reactive protein levels were associated with cardiovascular events in apparently healthy women, 22

Depressive symptoms were associated with increased mortality in older women, 78

Estrogen plus progestin was not effective for secondary prevention of coronary heart disease in postmenopausal women, 8

Long-term dietary folate was associated with a decreased risk for colon cancer in women, 51

Review: Postmenopausal hormone replacement therapy is associated with increased invasive ovarian cancer, especially after long-term use, 23

